

Due to the lapse in government funding, only web sites supporting excepted functions will be updated unless otherwise funded. As a result, the information on this website may not be up to date, the transactions submitted via the website may not be processed, and the agency may not be able to respond to inquiries until appropriations are enacted.

Updates regarding government operating status and resumption of normal operations can be found at <u>http://www.usa.gov</u>.

For Your Health: Recommendations for A Healthier You

Gay and bisexual men, like all men, need to be aware of the ways they can protect their health through all stages of life. For all men, the leading causes of death are heart disease and cancer. However, among men who have sex with men (MSM), there are higher rates of HIV and other sexually transmitted diseases (STDs), tobacco and drug use, and depression compared to other men.

<u>Sexual health (http://www.cdc.gov/sexualhealth/)</u>—the state of physical, emotional, mental, and social well being in relation to sexuality—is an important part of your overall health. The idea of sexual health emphasizes health and wellness as well as the avoidance of negative health outcomes that result from unhealthy sexual behaviors and attitudes. For gay and bisexual men, HIV, hepatitis, and other STDs are of particular concern. For example, the rate of new HIV diagnoses among MSM is more than 44 times that of other men, while the rate of primary and secondary syphilis among MSM is more than 46 times that of other men.



Many factors contribute to the higher rates of HIV and STDs among gay and bisexual men compared to the general population of men. These factors include high prevalence of HIV and other STDs among MSM, which increases the risk of disease exposure, and limits access to prevention services. Other factors are complacency about HIV risk, particularly among young gay and bisexual men; difficulty of consistently maintaining safe behaviors with every sexual encounter over the course of a lifetime; and lack of awareness of syphilis symptoms and how it can be transmitted (e.g., oral sex). Additionally, homophobia and stigma can prevent MSM from seeking and getting appropriate health services. Getting tested for HIV and STDs is a first step in protecting your health. Knowing you are infected allows you to get the care that you need to stay healthy and take steps to protect your partners.

What tests are recommended to help ensure the sexual health of gay and bisexual men?

As an individual, you can do a lot to protect your health. CDC recommends the following laboratory testing for gay and bisexual and other MSM:

- HIV (at least annually)
- Syphilis
- Chronic Hepatitis B infection
- **Hepatitis C** for men who engage in risky behaviors, such as rough sex or sex with multiple partners.
- Genital Herpes if directed by your health care provider.
- Chlamydia and Gonorrhea testing of the throat is needed if you have had receptive oral sex in the past year.
- **Chlamydia and Gonorrhea testing of the penis** (urethra) is needed if you have had insertive anal or oral sex in the past year.
- Chlamydia and Gonorrhea testing of the rectum is needed if you have had receptive anal sex in the past year.
- Visit <u>hivtest.org (http://www.hivtest.org)</u>
 [®] (http://www.cdc.gov/Other/disclaimer.html)</sup> or <u>findstdtest.org (http://www.findstdtest.org)</u>
 [®] (http://www.cdc.gov/Other /disclaimer.html)</sup> to finding testing locations.

You may want to get tested more often—every three to six months—if you have multiple or anonymous partners, have sex in conjunction with drug use, use methamphetamine, or have sex partners that participate in these activities.



What vaccinations does CDC recommend for gay and bisexual men?

- Hepatitis A and Hepatitis B vaccinations are recommended by CDC and the Advisory Committee on Immunization Practices (ACIP) because of higher rates of infection among gay and bisexual men.
 - $\circ~$ Two doses of the Hepatitis A vaccine are needed for lasting protection and the doses should be given at least six months apart.
 - A series of three or four doses of the Hepatitis B vaccine are usually given providing long-lasting protection.
- Seasonal flu and the H1N1 flu vaccinations are also recommended.
 - \circ Each vaccine is a single dose shot given before the start of the flu season in the fall.
- HPV vaccine is also available for gay and bisexual men up to 26 years of age to prevent genital warts and other <u>HPV-associated diseases and conditions (http://www.cdc.gov/hpv)</u>.
 - $\circ~$ The HPV vaccine is given as a three-dose series over a six month time period. It is best to be vaccinated before the first sexual contact, but later vaccination will protect those who have not been exposed to HPV.

How do I lower my risk for STDs?

There are three critical ways to protect yourself from STDs and HIV:

- Don't have sex (i.e. anal, vaginal, or oral)
- Be in a long-term, mutually monogamous sexual relationship with a partner you know has the same HIV status as you.
- Use a condom every time you have anal, vaginal, or oral sex. Consistent and correct use of the male latex condom reduces the risk of STD and HIV transmission. It is especially important to always use a condom during anal sex, since anal sex carries a much higher risk of HIV transmission.

It is also important to be honest and open with your doctor about your sexual behaviors so that he or she can give you the best and most appropriate care.

What other steps can I take to protect my health?

- Eat a healthy diet. Choosing healthful meal and snack options can help you avoid heart disease and its complications. Be sure to eat plenty of fresh fruits and vegetables.
- **Maintain a healthy weight.** To determine whether your weight is in a healthy range, doctors often calculate a number called the <u>body mass index (http://www.cdc.gov/healthyweight/assessing /bmi/)</u> (BMI). If you know your weight and height, you can calculate your BMI at CDC's <u>Assessing Your Weight (http://www.cdc.gov/healthyweight/assessing/index.html)</u> website. Or visit CDC's <u>Healthy Weight (http://www.cdc.gov/healthyweight/assessing/index.html)</u> website.
- **Exercise regularly.** Physical activity can help you maintain a healthy weight and lower cholesterol and blood pressure. Visit CDC's <u>Physical Activity (http://www.cdc.gov/physicalactivity /index.html)</u> site.
- **Don't smoke.** Cigarette smoking greatly increases your risk for heart disease, cancer, and stroke. So, if you don't smoke, don't start. If you do smoke, quitting will lower your risk for heart disease.
- Limit alcohol use. Avoid drinking too much alcohol, which can cause a variety of health problems (high blood pressure, cancer) and increase your risk of injury. Visit CDC's <u>Alcohol and Public Health (http://www.cdc.gov/alcohol/)</u> website.
- **Cholesterol screenings.** The National Cholesterol Education Program recommends that adults aged 20 years or older have their cholesterol checked every five years.
- **Cancer screenings.** Ask your health care provider for guidance on screening for prostate, testicular, colon, and anal cancers.
- **Check your blood pressure.** Getting your blood pressure checked is important because high blood pressure often has no symptoms.
- Get checkups. Ask your doctor or nurse how you can lower your chances for health problems.

How do I learn more?

- Talk to your health care provider.
- Call your local or state health department or contact an <u>LGBT Health Clinic</u> (<u>http://wwwdev.cdc.gov/lgbthealth/health-services.htm</u>).
- Contact CDC at 800-232-4636 (800-CDC-INFO) or cdcinfo@cdc.gov (mailto:cdcinfo@cdc.gov)

Related Information

- <u>Ten Things Gay Men Should Discuss with their Health Care Providers Gay & Lesbian Medical</u> <u>Association (http://www.glma.org/index.cfm?fuseaction=Page.viewPage&pageID=690)</u> (<u>http://www.cdc.gov/Other/disclaimer.html</u>)
- LGBT Health Clinics and Services (http://www.cdc.gov/lgbthealth/health-services.htm)
- <u>Medline Plus: Gay and Lesbian Health (http://www.nlm.nih.gov/medlineplus</u>

/gayandlesbianhealth.html) @ (http://www.cdc.gov/Other/disclaimer.html)

Page last reviewed: September 23, 2010 Page last updated: January 21, 2011 Content source: <u>National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention</u> Page maintained by: <u>National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention</u>

Centers for Disease Control and Prevention 1600 Clifton Rd. Atlanta, GA 30333, USA 800-CDC-INFO (800-232-4636) TTY: (888) 232-6348 - <u>Contact CDC–INFO</u>

